

References

- Astridge, D. J., Peeling, P., Goods, P. S. R., Girard, O., Hewlett, J., Rice, A. J., & Binnie, M. J. (2023). Rowing in Los Angeles: Performance Considerations for the Change to 1500 m at the 2028 Olympic Games. *Int J Sports Physiol Perform*, *18*(1), 104-107. <https://doi.org/10.1123/ijspp.2022-0231>
- de Campos Mello, F., de Moraes Bertuzzi, R. C., Grangeiro, P. M., & Franchini, E. (2009). Energy systems contributions in 2,000 m race simulation: a comparison among rowing ergometers and water. *European Journal of Applied Physiology*, *107*(5), 615-619. <https://doi.org/10.1007/s00421-009-1172-9>
- Fiskerstrand, Å., & Seiler, K. S. (2004). Training and performance characteristics among Norwegian International Rowers 1970–2001. *Scandinavian Journal of Medicine & Science in Sports*, *14*(5), 303-310. <https://doi.org/https://doi.org/10.1046/j.1600-0838.2003.370.x>
- Haugen, T., Sandbakk, Ø., Enoksen, E., Seiler, S., & Tønnessen, E. (2021). Crossing the Golden Training Divide: The Science and Practice of Training World-Class 800- and 1500-m Runners. *Sports Medicine*, *51*(9), 1835-1854. <https://doi.org/10.1007/s40279-021-01481-2>
- Maciejewski, H., Rahmani, A., Chorin, F., Lardy, J., Giroux, C., & Ratel, S. (2016). The 1,500-m Rowing Performance is Highly Dependent on Modified Wingate Anaerobic Test Performance in National-Level Adolescent Rowers. *Pediatr Exerc Sci*, *28*(4), 572-579. <https://doi.org/10.1123/pes.2015-0283>
- Manunzio, C., Mester, J., Kaiser, W., & Wahl, P. (2016). Training Intensity Distribution and Changes in Performance and Physiology of a 2nd Place Finisher Team of the Race across America Over a 6 Month Preparation Period [Original Research]. *Frontiers in Physiology*, *Volume 7 - 2016*. <https://doi.org/10.3389/fphys.2016.00642>
- Noordhof, D. A., Skiba, P. F., & de Koning, J. J. (2013). Determining anaerobic capacity in sporting activities. *Int J Sports Physiol Perform*, *8*(5), 475-482. <https://doi.org/10.1123/ijspp.8.5.475>
- Poole, D. C., Rossiter, H. B., Brooks, G. A., & Gladden, L. B. (2021). The anaerobic threshold: 50+ years of controversy. *J Physiol*, *599*(3), 737-767. <https://doi.org/10.1113/jp279963>
- Pripstein, L. P., Rhodes, E. C., McKenzie, D. C., & Coutts, K. D. (1999). Aerobic and anaerobic energy during a 2-km race simulation in female rowers. *European Journal of Applied Physiology and Occupational Physiology*, *79*(6), 491-494. <https://doi.org/10.1007/s004210050542>
- Rosenblat, M. A., Lin, E., da Costa, B. R., & Thomas, S. G. (2021). Programming Interval Training to Optimize Time-Trial Performance: A Systematic Review and Meta-

- Analysis. *Sports Med*, 51(8), 1687-1714. <https://doi.org/10.1007/s40279-021-01457-2>
- Rosenblat, M. A., Watt, J. A., Arnold, J. I., Treff, G., Sandbakk, Ø. B., Esteve-Lanao, J., Festa, L., Filipas, L., Galloway, S. D., Muñoz, I., Ramos-Campo, D. J., Schneeweiss, P., Sellés-Pérez, S., Stöggl, T., Talsnes, R. K., Zinner, C., & Seiler, S. (2025). Which Training Intensity Distribution Intervention will Produce the Greatest Improvements in Maximal Oxygen Uptake and Time-Trial Performance in Endurance Athletes? A Systematic Review and Network Meta-analysis of Individual Participant Data. *Sports Medicine*, 55(3), 655-673. <https://doi.org/10.1007/s40279-024-02149-3>
- Russell, A. P., Rossignol, P. F. L., & Sparrow, W. A. (1998). Prediction of elite schoolboy 2000-m rowing ergometer performance from metabolic, anthropometric and strength variables. *Journal of Sports Sciences*, 16(8), 749-754. <https://doi.org/10.1080/026404198366380>
- Sandford, G. N., Laursen, P. B., & Buchheit, M. (2021). Anaerobic Speed/Power Reserve and Sport Performance: Scientific Basis, Current Applications and Future Directions. *Sports Med*, 51(10), 2017-2028. <https://doi.org/10.1007/s40279-021-01523-9>
- Secher, N. H. (1993). Physiological and Biomechanical Aspects of Rowing. *Sports Medicine*, 15(1), 24-42. <https://doi.org/10.2165/00007256-199315010-00004>
- Seiler, K. S., & Kjerland, G. (2006). Quantifying training intensity distribution in elite endurance athletes: is there evidence for an "optimal" distribution? *Scand J Med Sci Sports*, 16(1), 49-56. <https://doi.org/10.1111/j.1600-0838.2004.00418.x>
- Seiler, S. (2010). What is best practice for training intensity and duration distribution in endurance athletes? *Int J Sports Physiol Perform*, 5(3), 276-291. <https://doi.org/10.1123/ijspp.5.3.276>
- Smith, T. B., & Hopkins, W. G. (2012). Measures of Rowing Performance. *Sports Medicine*, 42(4), 343-358. <https://doi.org/10.2165/11597230-000000000-00000>
- Sperlich, B., Matzka, M., & Holmberg, H.-C. (2023). The proportional distribution of training by elite endurance athletes at different intensities during different phases of the season [Review]. *Frontiers in Sports and Active Living*, Volume 5 - 2023. <https://doi.org/10.3389/fspor.2023.1258585>
- Stöggl, T., & Sperlich, B. (2014). Polarized training has greater impact on key endurance variables than threshold, high intensity, or high volume training [Original Research]. *Frontiers in Physiology*, Volume 5 - 2014. <https://doi.org/10.3389/fphys.2014.00033>
- Stöggl, T. L., & Sperlich, B. (2015). The training intensity distribution among well-trained and elite endurance athletes [Focused Review]. *Frontiers in Physiology*, Volume 6 - 2015. <https://doi.org/10.3389/fphys.2015.00295>

Zouhal, H., Le Douairon Lahaye, S., Abderrahaman, A. B., Minter, G., Herbez, R., & Castagna, C. (2012). Energy System Contribution to Olympic Distances in Flat Water Kayaking (500 and 1,000 m) in Highly Trained Subjects. *The Journal of Strength & Conditioning Research*, 26(3). https://journals.lww.com/nsca-jscr/fulltext/2012/03000/energy_system_contribution_to_olympic_distances_in.32.aspx